



Fertilization Program for:

“BANANA-PAPAYA”

RECOMMENDATIONS AND DOSAGES:

STAGE	PRODUCT	DOSAGE PER ACRE	APPLICATION
1 st Fertilization or Transplant	• Ergro [®] Set 20G	54 lb/acre	Application: 54 lb/acre of Ergro [®] Set 20G, to the soil mixed with the granular fertilizer, at the 1 st Fertilization. Banana: At the crop start. Papaya: At Transplant.
Growing	• Ergro [®] Set LSS	18 lb/acre	1 st Application: 9 lb/acre at Transplant (Papaya) or Crop Start (Banana). 2 nd Application: 9 lb/acre 30 days after the 1 st application. Via Drip Irrigation, Gravity or Drench.
Flowering	• Ergro [®] Crop • Ergro [®] Stim • Ergro [®] Win	½ quart/acre ½ quart/acre 4 lb/acre	1 st Foliar application of the 3 products at the beginning of flowering (Papaya) or cluster formation (Banana) (Ergro [®] Crop 8 fl oz/acre + Ergro [®] Stim 8 fl oz/acre + Ergro [®] Win 2 lb/acre). 2 nd Application 15 days after the 1 st application (Ergro [®] Crop 8 fl oz/acre + Ergro [®] Stim 8 fl oz/acre + Ergro [®] Win 2 lb/acre).
Maturity	• Ergro [®] Crop • Ergro [®] Stim • Ergro [®] K Excel	½ quart/acre ½ quart/acre 4 lb/acre	1 st Foliar application of the 3 products at the beginning of fruit formation (Ergro [®] Crop 8 fl oz/acre + Ergro [®] Stim 8 fl oz/acre + Ergro [®] K Excel 2 lb/acre). 2 nd Application 15 days after the 1 st application (Ergro [®] Crop 8 fl oz/acre + Ergro [®] Stim 8 fl oz/acre + Ergro [®] K Excel 2 lb/acre).