



Ergro® Crop

Recommendations and Dosage in Different Crops

Foliar Applications

Crop	Application Time	Dosage	Application
Vegetables Tomato Peppers Cucumber Eggplant Bróccoli Cauliflower Watermelon Cantaloupe Squash	Continuous Applications 1 st application 8 days after the transplant or after emerging. 2 nd application 15 days after the 1 st application. 3 rd We recommend applying at flowering stage repeat every 15 days and after the cut of fruits.	4-8 fl oz / acre 16 fl oz / acre 8 fl oz / acre	Foliar Application
Onions Garlic Potato Boniato Sweet Potato	Total of 3 applications 1 st application 15 days after emerging. 2 nd application 15 days after the 1 st application. 3 rd application 15 days after the 2 nd application.	1 st 8 a 24 fl oz / acre 2 nd 12 a 24 fl oz / acre 3 rd 16 a 32 fl oz / acre	Foliar Application
Cotton	Total of 2 applications 1 st application 15 days after emerging. 2 nd , 3 rd and 4 th applications 15 days after the previous one.	1 st 4 fl oz / acre 2 nd , 3 rd and 4 th 8 fl oz / acre	Foliar Application
Fruit Trees & Citrus Trees	High Portage Trees. Total of 3 applications 1 st application before flowering. 2 nd application during flowering. 3 rd application during at fructification. Low Portage Trees. Total of 3 applications 1 st application before flowering. 2 nd application during flowering. 3 rd application during at fructification.	1 st 16 a 24 fl oz / acre 2 nd 16 a 24 fl oz / acre 3 rd 24 a 32 fl oz / acre 1 st 8 - 16 fl oz / acre 2 nd 8 - 16 fl oz / acre 3 rd 8 - 16 fl oz / acre	Foliar Application Foliar Application
Basic Crops: Corn Sorghum Wheat Rice	From 2 to 3 applications. 1 st application 15 days after emerging. 2 nd application before flowering. 3 rd application before fructification.	1 st 8 fl oz / acre 2 nd 8 fl oz / acre 3 rd 16 fl oz / acre	Foliar Application
Beans Sor Beans	From 2 to 3 applications. 1 st application 8 days after emerging. 2 nd application during flowering. 3 rd application during at fructification.	1 st 4 fl oz / acre 2 nd 8 fl oz / acre 3 rd 8 fl oz / acre	Foliar Application
Sugarcane	Total of 3 applications. 1 st application 15 days after emerging. 2 nd application 30 days after the 1 st application. 3 rd application 30 days after the 2 nd application.	1 st 8 fl oz / acre 2 nd 16 fl oz / acre 3 rd 16 fl oz / acre	Foliar Application
Blueberry	Continuous Applications 1 st application beginning of the sprouting. 2 nd application to form the fruits. 3 rd application during at fructification.	1 st 4 fl oz / acre 2 nd 8 fl oz / acre 3 rd 8 fl oz / acre	Foliar Application
Floriculture	Continuous Applications Applied every 15 days.	4 a 8 fl oz / acre	Foliar Application