



Ergo[®] K Excel

Recommendations and Dosage in Different Crops

Foliar Applications

Crop	Application Time	Dosage	Application
Vegetables Tomato Pepper Cucumber Eggplant Squash	From 1 to 2 applications 1 st application: After the fruit formation. 2 nd application: 2 to 3 weeks after the first application.	1 st 2-2.5 lb / acre 2 nd 2-2.5 lb / acre	Foliar Application
Broccoli Cauliflower Watermelon Cantaloupe	From 1 to 2 applications 1 st application: After the fruit formation. 2 nd application: 2 to 3 weeks after the first application.	1 st 2-2.5 lb / acre 2 nd 2-2.5 lb / acre	Foliar Application
Cotton Garlic Onion Beans	From 1 to 2 applications 1 st application: After the fruit formation. 2 nd application: 2 to 3 weeks after the first application.	1 st 1.5-2 lb / acre 2 nd 1.5-2 lb / acre	Foliar Application
Corn Wheat Oat Barley Sorghum Saffron Rye Rice Chickpea	From 1 to 2 applications 1 st application: After the fruit formation. 2 nd application: 2 to 3 weeks after the first application.	1 st 1-2 lb / acre 2 nd 1-2 lb / acre	Foliar Application
Sugarcane	From 2 to 3 applications 1 st application: 3 months before harvest. 2 nd application: 2 months before harvest. 3 rd application: 1 month before harvest.	1 st 2-3 lb / acre 2 nd 2-3 lb / acre 3 rd 2-3 lb / acre	Foliar Application
Potato Sweet Potato	From 1 to 2 applications 1 st application: After the fruit formation. 2 nd application: 2 to 3 weeks after the first application.	1 st 2-3 lb / acre 2 nd 2-3 lb / acre	Foliar Application
Low Portage Fruit Trees	From 1 to 2 applications 1 st application: After the fruit formation. 2 nd application: 2 to 3 weeks after the first application.	1 st 2-2.5 lb / acre 2 nd 2-2.5 lb / acre	Foliar Application
High Portage Fruit Trees	From 1 to 2 applications 1 st application: After the fruit formation. 2 nd application: 2 to 3 weeks after the first application.	1 st 3-4 lb / acre 2 nd 3-4 lb / acre	Foliar Application