



Ergo[®] Amin L

Recommendations and Dosage in Different Crops

Foliar Applications

Crop	Application Time	Dosage	Application
Vegetables	Continuous Applications 1 st application one week after the transplant. 2 nd application 10 days after the 1 st application, every 10 to 15 days in each agrochemical application. 3 rd , 4 th , and subsequent 10 days after the 2 nd application.	1 st 4-8 fl oz / acre 2 nd 4-8 fl oz / acre 3 rd and 4 th 4-8 fl oz / acre	Foliar Application
Potato Sweet Potato Boniato	Total of 3 applications 1 st application 15 days after emerging. 2 nd application 15 days after the 1 st application. 3 rd application 15 days after the 2 nd application.	1 st 8 fl oz / acre 2 nd 12 fl oz / acre 3 rd 16 fl oz / acre	Foliar Application
Cotton	Total of 4 applications 1 st application at the beginning of flowering. 2 nd , 3 rd and 4 th applications 15 days after the previous one.	1 st 8 fl oz / acre 2 nd , 3 rd and 4 th 8 fl oz / acre	Foliar Application
Fruit Trees & Citrus Trees	High Portage Trees. Total of 3 applications 1 st application before flowering. 2 nd application during flowering. 3 rd application during at fructification and production.	High Portage Trees 1 st 16-24 fl oz / acre 2 nd 16-24 fl oz / acre 3 rd 16-24 fl oz / acre	Foliar Application
	Low Portage Trees. Total of 3 applications 1 st application before flowering. 2 nd application during flowering. 3 rd application during at fructification and production.	Low Portage Trees 1 st 8-16 fl oz / acre 2 nd 8-16 fl oz / acre 3 rd 8-16 fl oz / acre	Foliar Application
Basic Crops & Cereals	From 2 to 3 applications. 1 st application 20 days after emerging. 2 nd application, repeat before flowering. 3 rd application, repeat before fructification.	1 st 8-16 fl oz / acre 2 nd 8-16 fl oz / acre 3 rd 8-16 fl oz / acre	Foliar Application
Beans	From 2 to 3 applications. 1 st application before flowering. 2 nd application, repeat at flowering. 3 rd application, repeat before fructification.	1 st 4-8 fl oz / acre 2 nd 4-8 fl oz / acre 3 rd 4-8 fl oz / acre	Foliar Application
Sugarcane	Total of 3 applications. 1 st application 20-30 days after planting. 2 nd application 25-30 days after the 1 st application. 3 rd application 25-30 days after the 2 nd application.	1 st 8 fl oz / acre 2 nd 16 fl oz / acre 3 rd 16 fl oz / acre	Foliar Application
Berries	Continuous Applications 1 st Application at the beginning of flowering. 2 nd Application at the beginning of fruit formation. 3 rd , 4 th , 5 th , and subsequent applications every 15 days.	1 st 4 fl oz / acre 2 nd 8 fl oz / acre 3 rd and others 8 fl oz / acre	Foliar Application
Floriculture	Continuous Applications Apply every 2-3 weeks.	Each application of 8 fl oz in 30 gal of water. Approx. dosage per acre.	Foliar Application
Stress Crops	One or more Applications When the stress is present due to adverse conditions. Apply as many times as necessary with 1-2 weeks between each application.	Each application of 8 fl oz / acre.	Foliar Application

Soil Applications

All Crops	3 or More Applications 1 st application during growing. 2 nd application during flowering. 3 rd application during fructification.	Each application of 1-2 quarts / acre	Drip Irrigation, Channel Water, Micro Aspersions, Hydroponics, Semi- Hydroponics
------------------	--	---------------------------------------	--